

付表 4 痛み表現のクラスタごとの身体部位の平均選択率 (N=98-122)

クラスタ	身体部位															
	頭	歯	目	耳	のど	肩	背中	胸	腹	胃	腸	腰	手・腕	足・脚	関節	皮膚
A	.27	.06	.03	.03	.06	.05	.05	.26	<i>.19</i>	<i>.13</i>	.08	.06	.09	.07	.05	.05
B	<i>.18</i>	<i>.20</i>	.04	.08	.04	.04	.07	<i>.14</i>	<i>.17</i>	<i>.10</i>	.06	.09	<i>.12</i>	.10	<i>.16</i>	.06
C	.42	.24	.04	.04	.06	.03	.06	<i>.14</i>	<i>.18</i>	<i>.15</i>	.06	.05	.05	.05	.07	.05
D	.25	.27	.03	<i>.11</i>	.04	.04	.10	.08	.10	.05	.03	.06	<i>.13</i>	<i>.12</i>	<i>.10</i>	<i>.11</i>
E	.07	<i>.13</i>	<i>.18</i>	.06	.10	.04	.07	.09	.06	.05	.03	.06	<i>.18</i>	<i>.11</i>	.08	.25
F	<i>.12</i>	.09	.07	.03	<i>.17</i>	.07	.06	.21	<i>.13</i>	<i>.15</i>	.08	.06	<i>.13</i>	<i>.11</i>	.06	<i>.18</i>
G	.24	.06	.07	.05	.07	.21	.10	.07	<i>.11</i>	.09	.04	<i>.13</i>	.06	.09	.07	.03
H	<i>.18</i>	<i>.12</i>	<i>.12</i>	.04	<i>.12</i>	.07	.07	.10	<i>.13</i>	<i>.11</i>	.05	.08	.08	.08	.08	<i>.11</i>
I	.03	.04	.09	.07	<i>.17</i>	.03	.08	.03	.04	.06	.03	.04	.29	.31	<i>.11</i>	.26
J	<i>.14</i>	<i>.10</i>	.07	.03	<i>.10</i>	.06	.07	.09	<i>.11</i>	.08	.06	.06	<i>.18</i>	<i>.13</i>	.09	.24
K	.06	<i>.18</i>	.08	.08	.05	.06	.08	.09	.08	.07	.05	.07	<i>.15</i>	<i>.17</i>	<i>.16</i>	<i>.15</i>

A—K は Figure 4 のクラスタを示す。

太字は 20%以上の選択率を，斜体は 10%以上の選択率を示す。